



# PrimeVAC™

BY MERCK ANIMAL HEALTH

**Preconditioning programs for healthier,  
higher-value calves**



**MERCK**  
Animal Health

**The Science of Healthier Animals.™**



# Preconditioned to Perform

Proper preconditioning improves the efficiency and performance of calves, from the ranch to the feedyard. And that's why PrimeVAC™ by Merck Animal Health gives you a marketing edge. You can count on PrimeVAC preconditioning programs for healthier, higher-value calves that help you take advantage of value-added premiums.

Find the program that's right for your operation using trusted, reliable Merck Animal Health products. Talk to your veterinarian or Merck Animal Health representative about using PrimeVAC certification.

**PrimeVAC™**  
BY MERCK ANIMAL HEALTH  
CERTIFIED

Ranch Name: \_\_\_\_\_ Address: \_\_\_\_\_  
Contact Name: \_\_\_\_\_ City: \_\_\_\_\_  
Telephone: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_  
Email: \_\_\_\_\_  
No. of Head: \_\_\_\_\_ Breed: \_\_\_\_\_ Color: \_\_\_\_\_  
Gender:  Steer  Heifer  Mixed

The calves listed on this certificate have completed the health requirements for the following Merck Animal Health preconditioning program.

<p><b>PROGRAM</b></p> <p><input type="checkbox"/> PrimeVAC 24 <input type="checkbox"/> PrimeVAC 34 <input type="checkbox"/> PrimeVAC 34 Premium <input type="checkbox"/> PrimeVAC 45 <input type="checkbox"/> PrimeVAC 45 Premium <input type="checkbox"/> PrimeVAC Precon <input type="checkbox"/> PrimeVAC Heifer</p>	<p><b>PROTECTION</b></p> <p><b>Respiratory</b></p> <p><input type="checkbox"/> Vista™ Once SQ <input type="checkbox"/> Once PM8™ IN <input type="checkbox"/> Once PM8 SQ <input type="checkbox"/> Vista 5 SQ</p> <p><b>Reproductive</b></p> <p><input type="checkbox"/> Vista 5 VLS SQ</p> <p><b>Blockley</b></p> <p><input type="checkbox"/> Vision™ 7 <input type="checkbox"/> Vision 7 Somnus <input type="checkbox"/> Vision 8 <input type="checkbox"/> Vision 8 Somnus <input type="checkbox"/> Cavalry™ 9</p>	<p><b>CONTROL</b></p> <p><b>Internal Parasite</b></p> <p><input type="checkbox"/> Panacur™ Suspension <input type="checkbox"/> Safe-Guard™ Suspension</p> <p><b>External Parasite</b></p> <p><input type="checkbox"/> Ultra Boss™ Pour-On Insecticide <input type="checkbox"/> Ultra Saber™ Pour-On</p>
---	---	---

**PERFORMANCE**

**Implants**

Relegro™  
 Revalor™-G

Additional Notes: \_\_\_\_\_  
\_\_\_\_\_

<p><b>Certified by:</b> Owner Name: _____ Owner Signature: _____ Date: _____</p>	<p><b>Certified by:</b> Veterinarian Name: _____ Veterinarian Signature: _____ Date: _____</p>
--	--

This certificate is provided only as a service to the producer, owner or entity whose name appears on the face hereof ("Owner/Manager"). All information is derived from data supplied by the Owner/Manager, who is solely responsible for its accuracy and use. Merck Animal Health makes no warranties, express or implied, regarding the accuracy, adequacy, completeness, reliability, usefulness, or legality of any information. Such information is provided "as is", and should be verified by the user before relying upon it for any purpose.



Copyright © 2013 Intervet Inc., d/b/a Merck Animal Health, a subsidiary of Merck & Co., Inc.  
All rights reserved. 81-104-111-000-0002

Always consult your veterinarian concerning: best health management decisions specific to your operation, selection of qualified USDA/FDA approved products, optimum use of combination products, and the efficacy of vaccination in the face of maternal antibodies.

Always read, understand, and follow product label and use as directed.

Calves are vaccinated once:

- Administer vaccines at 3 to 4 months of age

## RESPIRATORY PROTECTION (ONE DOSE REQUIRED)

### IF DAMS ARE OPEN:

Vista<sup>®</sup> Once SQ

### IF DAMS ARE BRED:

Vista 5 SQ and either Once PMH<sup>®</sup> IN or Once PMH SQ

## BLACKLEG PROTECTION (ONE DOSE REQUIRED)

### SELECT ONE:

Vision<sup>®</sup> 7      Vision 8

Vision 7 Somnus      Vision 8 Somnus

Cavalry<sup>®</sup> 9

## INTERNAL PARASITE CONTROL (RECOMMENDED)

### SELECT ONE:

Safe-Guard<sup>®</sup> (fenbendazole)

Panacur<sup>®</sup> (fenbendazole)

## EXTERNAL PARASITE CONTROL (RECOMMENDED)

### SELECT ONE:

Ultra Boss<sup>®</sup> Pour-On Insecticide

Ultra Saber<sup>®</sup> Pour-On

## IMPLANT (OPTIONAL)

Ralgro<sup>®</sup> (zeranol)

# PrimeVAC 34

Calves are vaccinated twice:

- Administer the first vaccines at branding
- The second vaccination occurs 2 to 4 weeks prior to shipping

## RESPIRATORY PROTECTION (ONE DOSE REQUIRED)

**2 TO 4 WEEKS PRIOR TO SHIPPING:**

Vista 5 SQ and either Once PMH IN or Once PMH SQ

## BLACKLEG PROTECTION (TWO DOSES REQUIRED)

**SELECT ONE:**

Vision 7                      Vision 8  
Vision 7 Somnus          Vision 8 Somnus  
Cavalry 9

## INTERNAL PARASITE CONTROL (RECOMMENDED)

**2 TO 4 WEEKS PRIOR TO SHIPPING**

**SELECT ONE:**

Safe-Guard  
Panacur

## EXTERNAL PARASITE CONTROL (RECOMMENDED)

**2 TO 4 WEEKS PRIOR TO SHIPPING**

**SELECT ONE:**

Ultra Boss Pour-On Insecticide  
Ultra Saber Pour-On

## IMPLANT (OPTIONAL)

**AT BRANDING:**

Ralgro

# PrimeVAC 34 Premium

Calves are vaccinated twice:

- Administer the first vaccines at branding
- The second vaccination occurs 2 to 4 weeks prior to shipping

## RESPIRATORY PROTECTION (TWO DOSES REQUIRED)

**AT BRANDING:**

Vista 5 SQ

**2 TO 4 WEEKS PRIOR TO SHIPPING  
REVACCINATE WITH:**

Vista 5 SQ and either Once PMH IN or Once PMH SQ

## BLACKLEG PROTECTION (TWO DOSES REQUIRED)

**SELECT ONE:**

Vision 7

Vision 8

Vision 7 Somnus

Vision 8 Somnus

Cavalry 9

## INTERNAL PARASITE CONTROL (RECOMMENDED)

**2 TO 4 WEEKS PRIOR TO SHIPPING**

**SELECT ONE:**

Safe-Guard

Panacur

## EXTERNAL PARASITE CONTROL (RECOMMENDED)

**2 TO 4 WEEKS PRIOR TO SHIPPING**

**SELECT ONE:**

Ultra Boss Pour-On Insecticide

Ultra Saber Pour-On

## IMPLANT (OPTIONAL)

**AT BRANDING:**

Ralgro

# PrimeVAC 45

*For home-raised calves weaned a minimum of 45 days prior to delivery*  
**Calves are vaccinated twice:**

- Administer the first vaccines prior to or at weaning
- Revaccinate 3 to 6 weeks later

## **RESPIRATORY PROTECTION (TWO DOSES REQUIRED)**

### **IF FIRST VACCINATION AT PRE-WEANING:**

Vista 5 SQ and either Once PMH IN or Once PMH SQ  
and revaccinate with Vista 5 SQ

### **IF FIRST VACCINATION AT WEANING:**

Vista Once SQ and revaccinate with Vista 5 SQ

## **BLACKLEG PROTECTION (TWO DOSES REQUIRED)**

### **SELECT ONE:**

Vision 7                      Vision 8  
Vision 7 Somnus      Vision 8 Somnus  
Cavalry 9

## **INTERNAL PARASITE CONTROL (RECOMMENDED)**

### **AT PRE-WEANING OR WEANING PROCESSING**

#### **SELECT ONE:**

Safe-Guard  
Panacur

## **EXTERNAL PARASITE CONTROL (RECOMMENDED)**

### **AT PRE-WEANING OR WEANING PROCESSING**

#### **SELECT ONE:**

Ultra Boss Pour-On Insecticide  
Ultra Saber Pour-On

## **IMPLANT (OPTIONAL)**

### **AT BRANDING:**

Ralgro

# PrimeVAC 45 Premium

*For home-raised calves weaned a minimum of 45 days prior to delivery*

**Calves are vaccinated three times:**

- Administer the first vaccines at branding
- Administer the second vaccination prior to or at weaning
- Revaccinate 3 to 6 weeks later

## **RESPIRATORY PROTECTION (THREE DOSES REQUIRED)**

### **FIRST VACCINATION AT BRANDING:**

Vista 5 SQ and either Once PMH IN or Once PMH SQ

### **IF SECOND VACCINATION AT PRE-WEANING:**

Vista 5 SQ and either Once PMH IN or Once PMH SQ

### **IF SECOND VACCINATION AT WEANING:**

Vista Once SQ

### **THIRD VACCINATION 3 TO 6 WEEKS LATER:**

Vista 5 SQ

## **BLACKLEG PROTECTION (THREE DOSES REQUIRED)**

### **SELECT ONE:**

Vision 7                      Vision 8  
Vision 7 Somnus      Vision 8 Somnus  
Cavalry 9

## **INTERNAL PARASITE CONTROL (REQUIRED)**

### **AT PRE-WEANING OR WEANING PROCESSING**

#### **SELECT ONE:**

Safe-Guard  
Panacur

## **EXTERNAL PARASITE CONTROL (REQUIRED)**

### **AT PRE-WEANING OR WEANING PROCESSING**

#### **SELECT ONE:**

Ultra Boss Pour-On Insecticide  
Ultra Saber Pour-On

## **IMPLANT (OPTIONAL)**

### **AT BRANDING:**

Ralgro

### **AT WEANING:**

Ralgro or Revalor<sup>®</sup>-G

# PrimeVAC Precon

*For commingled calves weaned at least 60 days prior to delivery*

**Calves are vaccinated twice:**

- Administer the first vaccines within 24 hours of arrival
- Administer the second vaccination 3 to 6 weeks later and at least 14 days prior to shipping

## **RESPIRATORY PROTECTION (TWO DOSES REQUIRED)**

### **ON ARRIVAL:**

Vista Once SQ  
and revaccinate with Vista 5 SQ

## **BLACKLEG PROTECTION (TWO DOSES REQUIRED)**

### **SELECT ONE:**

Vision 7                      Vision 8  
Vision 7 Somnus          Vision 8 Somnus  
Cavalry 9

## **INTERNAL PARASITE CONTROL (RECOMMENDED)**

### **SELECT ONE:**

Safe-Guard  
Panacur

## **EXTERNAL PARASITE CONTROL (RECOMMENDED)**

### **SELECT ONE:**

Ultra Boss Pour-On Insecticide  
Ultra Saber Pour-On

## **IMPLANT (OPTIONAL)**

### **ON ARRIVAL**

### **SELECT ONE:**

Ralgro  
Revalor-G

# PrimeVAC Heifer

*For home-raised replacement heifers*

**Replacement heifers are vaccinated three times:**

- Administer first vaccines at pre-weaning or weaning
- Administer second vaccination 3 to 6 weeks later (including optional Brucellosis vaccination)
- Third vaccination 2 to 8 weeks prior to breeding

## **RESPIRATORY PROTECTION (TWO DOSES REQUIRED)**

### **IF FIRST VACCINATION AT PRE-WEANING:**

Vista 5 SQ and either Once PMH IN or Once PMH SQ  
and revaccinate with Vista 5 SQ

### **IF FIRST VACCINATION AT WEANING:**

Vista Once SQ  
and revaccinate with Vista 5 SQ

## **REPRODUCTIVE PROTECTION (ONE DOSE REQUIRED)**

### **2 TO 8 WEEKS PRIOR TO BREEDING:**

Vista 5 VL5 SQ

## **BLACKLEG PROTECTION (TWO DOSES REQUIRED)**

**PRIOR TO OR AT WEANING  
AND REVACCINATE 3 TO 6 WEEKS LATER  
SELECT ONE:**

Vision 7                      Cavalry 9                      Vision 8  
Vision 7 Somnus                      Vision 8 Somnus

## **INTERNAL PARASITE CONTROL (RECOMMENDED)**

**AT WEANING AND PRIOR TO BREEDING  
SELECT ONE:**

Safe-Guard  
Panacur

## **EXTERNAL PARASITE CONTROL (RECOMMENDED)**

**AT WEANING AND PRIOR TO BREEDING  
SELECT ONE:**

Ultra Boss Pour-On Insecticide  
Ultra Saber Pour-On

# Products

## RESPIRATORY PROTECTION

Only Vista vaccines offer the most complete dual-action pneumonia protection.



10 dose  
50 dose

### Vista Once SQ

Vista Once SQ combines the protection of Vista 5 SQ and Once PMH SQ. It is a 5-way MLV vaccine in combination with avirulent-live *Pasteurella multocida* and *Mannheimia haemolytica*.

Protection against:

- *Mannheimia haemolytica*
- *Pasteurella multocida*
- IBR (Infectious Bovine Rhinotracheitis)
- BVD Type 1 (Bovine Viral Diarrhea Type 1)
- BVD Type 2 (Bovine Viral Diarrhea Type 2)
- PI<sub>3</sub> (Parainfluenza<sub>3</sub>)
- BRSV (Bovine Respiratory Syncytial Virus)



10 dose  
50 dose

### Vista 5 SQ

5-way MLV vaccine.

Protection against:

- IBR (Infectious Bovine Rhinotracheitis)
- BVD Type 1 (Bovine Viral Diarrhea Type 1)
- BVD Type 2 (Bovine Viral Diarrhea Type 2)
- PI<sub>3</sub> (Parainfluenza<sub>3</sub>)
- BRSV (Bovine Respiratory Syncytial Virus)



10 dose  
50 dose

## Once PMH IN

The first and only intranasal vaccine against bacterial pneumonia.

Protection against:

- *Mannheimia haemolytica*
- *Pasteurella multocida*

Approved for all ages, including calves 1 week old.



10 dose  
50 dose

## Once PMH SQ

The only avirulent-live dual-action bacterial pneumonia vaccine available.

Protection against:

- *Mannheimia haemolytica*
- *Pasteurella multocida*

# Products

## BLACKLEG PROTECTION

Tested and trusted clostridial vaccines. Vision vaccines are proven to be the smoothest clostridial vaccines on the market, with less stress and fewer injection site reactions.



### Vision 7

7-way clostridial protection:

- *Clostridium chauvoei* (Blackleg)
  - *Clostridium septicum* (Malignant Edema)
  - *Clostridium novyi* (Black Disease)
  - *Clostridium sordelli*
  - *Clostridium perfringens* Types C and D (Overeating)
- 2mL subcutaneous injection
  - Available in 10, 50 and 250 doses

### Vision 7 Somnus

7-Way Clostridial/Somnus Protection:

- *Clostridium chauvoei* (Blackleg)
  - *Clostridium septicum* (Malignant Edema)
  - *Clostridium novyi* (Black Disease)
  - *Clostridium sordelli*
  - *Clostridium perfringens* Types C and D (Overeating)
  - *Haemophilus somnus* (*Histophilus somni*)
- The only low-dose, 7-way clostridial/somnus combination available
  - 2mL subcutaneous injection
  - Available in 10, 50 and 250 dose

## Vision 8

8-way clostridial protection:

- *Clostridium haemolyticum* (Redwater)
  - *Clostridium chauvoei* (Blackleg)
  - *Clostridium septicum* (Malignant Edema)
  - *Clostridium novyi* (Black Disease)
  - *Clostridium sordelli*
  - *Clostridium perfringens* Types C and D (Overeating)
- 2mL subcutaneous injection
  - Available in 10 and 50 doses

## Vision 8 Somnus

8-way clostridial protection:

- *Clostridium haemolyticum* (Redwater)
  - *Clostridium chauvoei* (Blackleg)
  - *Clostridium septicum* (Malignant Edema)
  - *Clostridium novyi* (Black Disease)
  - *Clostridium sordelli*
  - *Clostridium perfringens* Types C and D (Overeating)
  - *Haemophilus somnus* (*Histophilus somni*)
- The only low-dose, 8-way clostridial/somnus combination available
  - 2mL subcutaneous injection
  - Available in 10 and 50 doses

## Cavalry 9

The only 9-way clostridial protection including tetanus:

- *Clostridium chauvoei* (Blackleg)
  - *Clostridium septicum* (Malignant Edema)
  - *Clostridium novyi* Type B (Black Disease)
  - *Clostridium haemolyticum* (Redwater)
  - *Clostridium sordelli*
  - *Clostridium perfringens* Types C and D (Overeating)
  - *Clostridium tetani* (Tetanus)
- Available in 10, 50 and 125 doses

# Products

## INTERNAL PARASITE CONTROL

Safe and proven parasite control.



### **Panacur** (fenbendazole) **Suspension** (Drench)

- Available from your veterinarian
- For removal and control of lungworms, stomach worms and intestinal worms
- Easy-to-use applicator gun for accurate and stress-free deworming
- One liter deworms 43,478 lb. of cattle
- One gallon deworms 164,565 lb. of cattle
- 2.33cc/100 lb. of cattle

### **Safe-Guard** (fenbendazole) **Suspension** (Drench)

- Single application
- Low-dose volume
- Easy-to-use applicator gun for accurate and stress-free deworming
- One liter deworms 43,478 lb. of cattle
- One gallon deworms 164,565 lb. of cattle
- 2.33cc/100 lb. of cattle

Panacur (fenbendazole) Suspension Drench & Safe-Guard (fenbendazole) Drench and Paste: **RESIDUE WARNING:** Cattle must not be slaughtered within eight days following last treatment. For dairy cattle, the milk discard time is zero hours. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

# Panacur®

(fenbendazole)

## Horse & Cattle Dewormer

### Suspension 10% (100 mg/mL)

EXAMPLES: (Horses and Cattle)

Dose (5 mg / kg)	Dose (10 mg / kg)	Animal Weight
2.3 mL	4.6 mL	100 lb
4.6 mL	9.2 mL	200 lb
6.9 mL	13.8 mL	300 lb
9.2 mL	18.4 mL	400 lb
11.5 mL	23.0 mL	500 lb
23.0 mL	46.0 mL	1,000 lb
34.5 mL	69.0 mL	1,500 lb

#### ► RESIDUE WARNINGS: ◀

- Do not use in horses intended for food.
- Cattle must not be slaughtered for human consumption within 8 days following treatment.
- Do not use at 10 mg/kg in dairy cattle. Dose rate of 10 mg/kg is for beef cattle only. Dose rate of 10 mg/kg in dairy cattle could result in violative residues in milk.
- A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Keep this and all medication out of the reach of children.

**1,000 mL (33.8 fl.oz)**

10 mg/kg (4.6 mg/lb) for the control of ascarids (*Parascaris equorum*). EXAMPLE: (10 mg/kg); 2.3 mL/50 lb; 23 mL/500 lb

**Beef and Dairy Cattle:** 5 mg/kg (2.3 mg/lb) for the removal and control of:

**Lungworm:** (*Dictyocaulus viviparus*).

**Stomach worm (adults):** *Ostertagia ostertagi* (brown stomach worm). **Stomach worm (adults & 4th stage larvae):** *Haemonchus contortus/pla- ceii* (barberpole worm), *Trichostrongylus axei* (small stomach worm).

**Intestinal worm (adults & 4th stage larvae):** *Bunostomum phlebotomum* (hookworm), *Nematodirus helvetianus* (thread-necked intestinal worm), *Cooperia punctata* and *C. oncophora* (small intestinal worm), *Trichostrongylus colubri- formis* (bankrupt worm), *Oesophagostomum radiatum* (nodular worm).

**Beef Cattle Only - 10 mg/kg (4.6 mg/lb) for the removal and control of:**

**Stomach worm (4th stage inhibited larvae):** *Ostertagia ostertagi* (Type II Ostertagiasis).

**Tapeworm:** *Moniezia benedeni*.

#### ► Do not use in dairy cattle at 10 mg/kg ◀

In beef and dairy cattle, the recommended dose of 5 mg/kg is achieved when 2.3 mL of the drug are given for each 100 lb of body weight. In beef cattle only, the recommended dosage of 10 mg/kg for the treatment of Ostertagiasis Type II

#### DOSAGE:

**Horses:** 5 mg/kg (2.3 mg/lb) for the control of large strongyles (*Strongylus edentatus*, *S. equi- nus*, *S. vulgaris*, *Triodontophorus* spp.), small strongyles (*Cyathostomum* spp., *Cylicocyclus* spp., *Cylicostephanus* spp., *Cylicodontophorus* spp.), and pinworms (*Oxyuris equi*). EXAMPLE: 2.3 mL/100 lb; 23 mL/1,000 lb

(inhibited 4th stage larvae) or tapeworm is achieved when 4.6 mL of the drug is given for each 100 lb of body weight.

#### DIRECTIONS:

**Beef and Dairy Cattle and Horses:** Determine the proper dose according to estimated body weight. Administer orally by suitable dosing syringe. Insert nozzle of syringe through the interdental space and deposit the drug on the back of the tongue by depressing the plunger. The drug may also be administered by stomach tube. There are no known contraindications to the use of the drug in cattle or horses. For dairy cattle, there is no milk withdrawal period at 5 mg/kg. Panacur® (fenbendazole) Suspension 10% is approved for use concomitantly with an approved form of trichlorfon. Trichlorfon is approved for the treatment of stomach bots (*Gasterophilus* spp.) in horses. Refer to the manufacturer's label for directions for use and cautions for trichlorfon.

Regular deworming at intervals of six to eight weeks may be required for horses.

Under conditions of continued exposure to parasites, retreatment of cattle may be needed after 4-6 weeks.

697700-B

# Products

## EXTERNAL PARASITE CONTROL

Fast-acting, long-lasting insect control.



### Ultra-Boss Pour-On

5% Permethrin, 5% Piperonyl Butoxide

- Kills and controls the face flies that spread pinkeye
- Effective against face flies, horn flies, lice, horse flies, stable flies, mosquitoes, back flies and ticks
- Convenient low dose:
  - 10mL for animals 600 lb. or less
  - 15mL for animals over 600 lb.

### Ultra Saber Pour-On

1% Lambda-cyhalothrin

- Long-lasting control of horn flies and lice
- Convenient low dose
  - 10mL for animals 600 lb. or less
  - 15mL for animals over 600 lb.

## IMPLANTS

Reliable, safe and effective implants for cattle on grass.



### Ralgro

- Approved to increase rate of weight gain in suckling calves
- Increases rate of weight gain and feed conversion of weaned beef calves, growing beef cattle, feedlot steers and heifers
- Stimulates animal's own natural growth system
- Shown to increase weaning weight by as much as 20-35 lb. in calves

*A withdrawal period has not been established for Ralgro in pre-ruminating calves. Do not use in calves to be processed for veal. For complete information, refer to product label.*



### Revalor-G

40 mg of trenbolone acetate and 8 mg of estradiol

- Increase rate of weight gain in weaned pasture steers and heifers
- Uses a lower hormone dosage, designed for stocker cattle on pasture
- Implanting gun available

## REPRODUCTIVE PROTECTION

Only Vista vaccines offer single-dose fetal protection against BVD for the entire pregnancy.



10 dose  
50 dose

### Vista 5 VL5 SQ

5-way modified-live virus vaccine with *Campylobacter fetus* (vibriosis) and 5-way Leptospirosis protection.

Modified-live virus vaccine, protects against:

- IBR (Infectious Bovine Rhinotracheitis)
- BVD Type 1 (Bovine Viral Diarrhea Type 1)
- BVD Type 2 (Bovine Viral Diarrhea Type 2)
- PI<sub>3</sub> (Parainfluenza<sub>3</sub>)
- BRSV (Bovine Respiratory Syncytial Virus)
- 5 strains of Leptospirosis, including *L. hardjo-ovis*
- *Campylobacter fetus* (vibriosis) protection





# YOUR LIVELIHOOD IS OUR RESPONSIBILITY.

On ranches, dairies and feedyards across America, the science of healthier animals is a way of life. It's this responsibility that drives who we are and what we do.

We're Merck Animal Health, and we're dedicated to improving the health and well-being of animals through innovative, science-based products, treatments and services.

Because of your efforts, we all enjoy a dependable, affordable food supply. You have our thanks and support. Please contact your veterinarian, animal health supplier or Merck Animal Health representative for more information.

We're with you every step of the way.



**The Science of Healthier Animals.™**

Visit [merck-animal-health-usa.com](http://merck-animal-health-usa.com)